


Cancer Risk after Fat Transfer: A Multicenter Case-Cohort Study

Terence M. Myckatyn, M.D.
 I. Janelle Wagner, M.D.
 Babak J. Mehrara, M.D.
 Melissa A. Crosby, M.D.
 Julie E. Park, M.D.
 Bahjat F. Qaqish, M.D.,
 Ph.D.
 Dominic T. Moore, M.P.H.
 Evan L. Busch, Ph.D.
 Amanda K. Silva, M.D.
 Surinder Kaur, Ph.D.
 David W. Ollila, M.D.
 Clara N. Lee, M.D., M.P.P.

St. Louis, Mo.; Chapel Hill, N.C.; New York, N.Y.; Houston, Texas; Chicago and Arlington Heights, Ill.; Boston, Mass.; and Columbus, Ohio



Background: Fat transfer is an increasingly popular method for refining post-mastectomy breast reconstructions. However, concern persists that fat transfer may promote disease recurrence. Adipocytes are derived from adipose-derived stem cells and express adipocytokines that can facilitate active breast cancer cells in laboratory models. The authors sought to evaluate the association between fat transfer to the reconstructed breast and cancer recurrence in patients diagnosed with local or regional invasive breast cancers.

Methods: A multicenter, case-cohort study was performed. Eligible patients from four centers (Memorial Sloan Kettering, M. D. Anderson Cancer Center, Alvin J. Siteman Cancer Center, and the University of Chicago) were identified by each site's institutional tumor registry or cancer data warehouse. Eligibility criteria were as follows: mastectomy with immediate breast reconstruction between 2006 and 2011, age older than 21 years, female sex, and incident diagnosis of invasive ductal carcinoma (stage I, II, or III). Cases consisted of all recurrences during the study period, and controls consisted of a 30 percent random sample of the study population. Cox proportional hazards regression was used to evaluate for association between fat transfer and time to recurrence in bivariate and multivariate models.

Results: The time to disease recurrence unadjusted hazard ratio for fat transfer was 0.99 (95 percent CI, 0.56 to 1.7). After adjustment for age, body mass index, stage, HER2/Neu receptor status, and estrogen receptor status, the hazard ratio was 0.97 (95 percent CI, 0.54 to 1.8).

Conclusion: In this population of breast cancer patients who had mastectomy with immediate reconstruction, fat transfer was not associated with a higher risk of cancer recurrence. (*Plast. Reconstr. Surg.* 139: 11, 2017.)

CLINICAL QUESTION/LEVEL OF EVIDENCE: Therapeutic, III.

Fat transfer has gained widespread acceptance as a surgical technique for volume restoration and contour correction in breast reconstruction.¹ Members of the American Society of Plastic Surgeons report performing 25,456

fat transfer procedures in 2014,² and 62 percent of surveyed members use this technique for breast reconstruction.³ Despite its utility in improving

From the Division of Plastic and Reconstructive Surgery, Washington University School of Medicine in Saint Louis, and the Alvin J. Siteman Cancer Center; the Division of Plastic and Reconstructive Surgery, the Department of Biostatistics, the Department of Epidemiology, Gillings School of Global Public Health, the Lineberger Comprehensive Cancer Center, and the Division of Surgical Oncology, Department of Surgery, University of North Carolina; the Department of Surgery, Memorial Sloan Kettering Cancer Center; the Department of Plastic Surgery, M. D. Anderson Cancer Center; the Section of Plastic and Reconstructive Surgery, University of Chicago Medicine and Biological Sciences, University of Chicago; The Plastic Surgery Foundation; the Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School; the Department of Epidemiology, Harvard T. H. Chan School of Public Health; and Plastic and Reconstructive Surgery, Health Services Management and Policy, Ohio State University.

Received for publication April 4, 2016; accepted August 4, 2016.

Copyright © 2016 by the American Society of Plastic Surgeons

DOI: 10.1097/PRS.0000000000002838

Disclosure: The authors have no financial interest to declare in relation to the content of this article.

A "Hot Topic Video" by Editor-in-Chief Rod J. Rohrich, M.D., accompanies this article. Go to PRSJJournal.com and click on "Plastic Surgery Hot Topics" in the "Videos" tab to watch. On the iPad, tap on the Hot Topics icon.



This work was supported by
THE PLASTIC SURGERY FOUNDATION.

aesthetic outcomes and optimization of symmetry following mastectomy with reconstruction, concerns regarding its oncologic safety persist.⁴⁻⁹ These concerns are based on laboratory studies demonstrating that adipose-derived stem cells and adipose-derived growth factors can modulate the behavior of breast tumors *in vitro* and in animal models.¹⁰⁻¹⁸ Also, laboratory studies have shown that adipose-derived stem cells modulate desmoplasia by elaborating extracellular matrix proteins, attenuate the antitumor immune response, and promote angiogenesis.^{4,5,19,20} A few retrospective clinical studies have suggested that fat transfer may increase the risk of locoregional recurrence after mastectomy for ductal carcinoma *in situ* or following partial mastectomy.^{6-8,21} Aside from a recent matched controlled study that shows fat transfer to be oncologically safe,²² most clinical studies have been limited by inadequate power to detect small effects.

Recent guiding principles published by the American Society of Plastic Surgeons acknowledge that a limited body of evidence shows fat transfer following postmastectomy breast reconstruction to be oncologically safe.²³ These guiding principles, however, also acknowledge the need for additional high-quality studies. As such, the American Society of Plastic Surgeons clinical trials committee sought to establish whether adjunctive fat transfer is associated with a higher risk of recurrence in patients who have undergone mastectomy with reconstruction for invasive breast cancer. Our experimental design took into consideration the relatively low baseline rate of cancer recurrence, and the fact that although fat transfer is very popular currently, it gained prominence as a technique relatively recently. Moreover, we recognized the immediate need for information examining the impact of fat transfer on cancer recurrence given its popularity—something that a prospective trial could not provide. The design of this study improves on previous work, with more representative selection of controls, adjustment for duration of follow-up, and sufficient power to detect a doubling of breast cancer recurrence risk.

PATIENTS AND METHODS

Study Design

A case-cohort design was used. The case-cohort approach allows for greater precision in the circumstance of a rare outcome and adjustment for different durations of follow-up.

Study Population

Patients were identified through the tumor registry or data warehouse of four sites: University of Chicago, M. D. Anderson Cancer Center, Memorial Sloan Kettering Cancer Center, and the Siteman Cancer Center at Washington University School of Medicine in St. Louis. Institutional review board approval was obtained at each site. Eligible patients consisted of all women 21 years and older with incident invasive ductal carcinoma, stages I through III, who were diagnosed between January 1, 2006, and December 31, 2011, and treated with mastectomy and immediate breast reconstruction. We excluded men, women younger than 21 years, women with prior breast cancer, and women with stage IV or inflammatory breast cancer. We also excluded women who had delayed breast reconstruction to minimize heterogeneity in time intervals between diagnosis, treatment, and fat transfer. Only patients with invasive ductal carcinoma were included, so patients with ductal carcinoma *in situ* only, lobular carcinoma *in situ* only, sarcoma, invasive lobular carcinoma, or no cancer were excluded.

Identification of Cases and Controls

Cases consisted of all eligible patients who had experienced a recurrence (local, regional, or distant) during the study period (January 1, 2006, to December 31, 2011) as reported by each site's tumor registry or cancer data warehouse. The cohort was a 30 percent random sample of the control population, defined as patients who did not have a recurrence during the study period. Exposure to fat grafting was measured using medical record review in three sites and a prospectively maintained plastic surgery clinical database (at Memorial Sloan Kettering Cancer Center). The approach of Cai and Zeng was used for power and sample size considerations.²⁴

Statistical Analyses

Fisher's exact tests were used to evaluate general association for categorical data. The Wilcoxon rank sum test (using Van der Waerden or normal scores) was used for two-group comparisons of continuous covariates. Cox proportional hazards regression modeling was used to explore the association of covariates of interest with time to recurrence. Time to recurrence was defined as the time from the date of diagnosis to the date of a local, regional, or distant cancer recurrence. Patients who reached the end of the study period, were lost to follow-up, or died without documented disease

recurrence were considered censored. The covariates of interest were as follows: fat transfer (yes or no), age, tumor stage (I through III), smoking status (yes or no), body mass index, estrogen receptor status (positive or negative), progesterone receptor status (positive or negative), HER2/Neu amplification status (positive or negative), receipt of adjuvant chemotherapy (yes or no), receipt of neoadjuvant chemotherapy (yes or no), receipt of adjuvant radiation therapy (yes or no), and receipt of adjuvant endocrine therapy (yes or no).

We report both bivariate and multivariate estimated hazard ratios with 95 percent confidence intervals. The final multivariate models did not include progesterone receptor status because it was highly correlated with estrogen receptor status. They did not include adjuvant therapies because they were highly correlated with stage. Statistical analyses were performed using SAS (Version 9.3; SAS Institute, Inc., Cary, N.C.) and R.²⁵

RESULTS

We identified 3271 eligible patients across the four institutions. The study sample ($n = 1197$) consisted of all recurrences during the study period ($n = 225$) and a 30 percent random sample of the study population ($n = 972$) (Table 1). Based on this sample size, power was calculated to be 76 percent against a relative risk of 2 and 86 percent against a relative risk of 2.22, when using a one-sided test with type I error of 5 percent.

The median age was 47 ± 10.1 years. The median body mass index was 26.5 ± 5.6 kg/m². Detailed in Table 1, almost half of the patients in the study sample had stage I disease, and most had estrogen receptor–positive and progesterone receptor–positive tumors. More than half received adjuvant chemotherapy, and approximately one-quarter received radiation therapy, endocrine therapy, or neoadjuvant chemotherapy. Although 80 percent of patients underwent reconstruction with breast implants, the remainder underwent reconstruction with autologous flaps, or a combination of flap and implant. Fat transfer was performed in 64 patients (5 percent), including 28 at M. D. Anderson, 26 at Memorial Sloan Kettering, seven at Washington University, and three at the University of Chicago.

Two hundred twenty-five patients, or 6.9 percent of the entire study population, had a recurrence of breast cancer. Of these, 124 recurrences were distant (55 percent), 24 were regional (11 percent), and 77 were local (34 percent). Forty-eight patients (4 percent) died during the study period. In bivariate analyses of associations between individual covariates of interest and time to recurrence (Table 2), patients who underwent fat transfer had an equivalent risk of cancer recurrence relative to those who did not (hazard ratio, 0.99; 95 percent CI, 0.56 to 1.7). Patients with HER2/Neu-positive tumors had a lower hazard of cancer recurrence than patients with HER2/Neu-negative tumors (hazard ratio, 0.62; 95 percent CI, 0.42 to 0.91). Patients with estrogen

Table 1. Patient Characteristics

Patient Characteristics	Total Cohort*	Recurrence	No Recurrence
Mean age (SD), yr	47.78 (10.09)	44	48
Mean BMI (SD), kg/m ²	26.49 (5.62)	25	25
TNM stage			
I	585 (49)	77 (34)	508 (52)
II	448 (37)	92 (41)	356 (37)
III	164 (14)	56 (25)	108 (11)
Receptors			
HER2Neu ⁺	215 (20)	29 (13)	205 (22)
ER ⁺	875 (74)	134 (60)	741 (78)
PR ⁺	713 (61)	101 (46)	612 (64)
Treatment			
Adjuvant radiation therapy	319 (27)	110 (49)	209 (22)
Adjuvant chemotherapy	689 (58)	153 (68)	536 (55)
Adjuvant endocrine therapy	1 (0)	25 (42)	252 (74)
Neoadjuvant chemotherapy	250 (21)	82 (36)	168 (17)
Site			
M. D. Anderson	271 (23)	32 (14)	239 (25)
Memorial Sloan Kettering	794 (66)	166 (74)	628 (65)
University of Chicago	39 (3)	8 (4)	31 (3)
Washington University in St. Louis	93 (8)	19 (8)	74 (8)
Totals	1197 (100)	225 (19)	972 (81)

BMI, body mass index; TNM, tumor, node, metastasis; ER, estrogen receptor; PR, progesterone receptor.

*Values in parentheses for age and BMI represent standard deviation. All other values in parentheses represent a percentage.

Table 2. Bivariate Time to Recurrence Models

Covariates	Hazard Ratio	95% CI	<i>p</i>
Age (for each 10 yr)	0.81	0.71–0.93	0.002
BMI (kg/m ²)	0.98	0.96–1.01	0.19
Stage II vs. stage I	1.5	1.1–2.1	0.007
Stage III vs. stage II	2	1.4–2.8	<0.0001
HER2Neu ⁺ vs. HER2Neu ⁻	0.62	0.42–0.91	0.02
ER ⁺ vs. ER ⁻	0.51	0.39–0.66	<0.0001
PR ⁺ vs. PR ⁻	0.52	0.40–0.68	<0.0001
Fat transfer (yes vs. no)	0.99	0.56–1.7	0.99

BMI, body mass index; ER, estrogen receptor; PR, progesterone receptor.

receptor–positive tumors had 51 percent of the risk of patients with estrogen receptor–negative tumors ($p < 0.0001$). Stage II patients had a greater hazard of cancer recurrence than stage I patients (hazard ratio, 1.5; 95 percent CI, 1.1 to 2.1) and stage III patients had a greater hazard of cancer recurrence than stage II patients (hazard ratio, 2.0; 95 percent CI, 1.4 to 2.8). In the multivariable model, adjusting for age, stage, body mass index, HER2/Neu-positive, and estrogen receptor–positive, patients who had fat transfer had a similar risk of recurrence as those who did not have fat transfer (risk, 97 percent; $p = 0.93$) (Table 3).

DISCUSSION

Fat transfer was not associated with an increased probability of breast cancer recurrence in this multicenter case-cohort study. Although fat transfer to the breast has also been used in the context of reconstruction for partial mastectomy,²⁶ or as the sole technique of breast reconstruction following mastectomy,²⁷ we investigated fat transfer as an adjunctive technique to prosthetic or flap-based reconstruction. Our findings are primarily applicable for patients with stage I through III invasive ductal carcinomas treated with mastectomy and immediate breast reconstruction.

Although several epidemiologic studies report a link between obesity in postmenopausal women and breast cancer,^{28–31} and translational research studies report that mesenchymal cells or

adipose-derived stem cells support the progression of existing tumors,^{10,15,16,32–35} none shows that adipocytes form tumors de novo.⁵ As an endocrine organ, white adipose tissue may promote breast cancer through the secretion of adipokines such as leptin,^{36,37} or insulin-like growth factor.³⁸ Furthermore, reduced levels of adiponectin in obese patients fosters a permissive environment for the pro-oncogenic properties of leptin.^{37,39}

Adipose-derived stem cells offer another mechanism by which white adipose tissue can simulate breast cancer cells. The progression of breast cancer is impacted by stromal cells of mesenchymal and hematopoietic origin.⁵ Under defined conditions, adipocytes⁴⁰ and their progenitors promote tumorigenesis in both in vivo and in vitro models.⁴¹ When adipose-derived stem cells are exposed to tumor-conditioned media secreted by breast cancer cell lines, they tend to proliferate, differentiate into myofibroblasts, enhance tissue stiffness through altered extracellular matrix deposition, secrete proangiogenic factors, and exhibit attenuated adipogenic differentiation.⁴¹ Adipose-derived stem cells preferentially contribute vascular and fibrovascular tumor-associated fibroblasts to the tumor stroma, and bone marrow–derived mesenchymal stem cells contribute fibroblast-specific proteins.³⁴ Co-culture of adipose-derived stem cells with breast cancer cells can also facilitate tumor metastases.^{15,42,43} These important translational data speak to the potential consequences of using purified adipose-derived stem cell grafts in the presence of active cancer cells. They may lead to the identification of molecular markers to predict cancer recurrence after fat grafting⁴² but do not necessarily translate directly to current clinical practice. Unlike immunocompromised nude mice receiving purified adipose-derived stem cell cultures,⁴³ immunocompetent human patients typically receive fat grafts containing a variable but low (2 to 8 percent) fraction of adipose-derived stem cells.^{44,45}

The absence of an association between fat transfer and recurrence risk reported here is consistent with experimental studies showing that white adipose tissue may stimulate active but not dormant breast cancers. Human adipose-derived stem cells significantly increase their malignant potential when co-cultured with active, but not dormant, breast cancer cells in vitro and in a xenogenic murine recipient in vivo model.¹⁵ Under experimental conditions, invasive breast cancer cells alter the phenotype of adjacent mature

Table 3. Multivariate Time to Recurrence Model

Covariates	Hazard Ratio	95% CI	<i>p</i>
Age (for each 10 yr)	0.85	0.75–0.98	0.02
BMI (kg/m ²)	0.98	0.96–1.01	0.11
Stage II vs. stage I	1.3	0.95–1.8	0.13
Stage III vs. stage II	2.3	1.6–3.2	<0.0001
HER2Neu ⁺ vs. HER2Neu ⁻	0.48	0.32–0.72	0.0004
ER ⁺ vs. ER ⁻	0.48	0.37–0.63	<0.0001
Fat transfer (yes vs. no)	0.97	0.54–1.8	0.93

BMI, body mass index; ER, estrogen receptor; PR, progesterone receptor.

adipocytes which, in turn, may increase the malignant potential of local breast cancer cells.⁴⁰

Our findings are applicable to patients undergoing fat transfer following prosthetic or autologous reconstruction of mastectomy defects in the setting of stage I through III breast cancer. Following mastectomy, the engrafted recipient site contains little to no breast tissue. Although some studies have suggested that adipocytes or their progenitors may stimulate active breast cancer,^{10,16} this could be less problematic if susceptible tissues have already been extirpated. By contrast, a significant percentage of residual breast tissue remains following partial mastectomy, another context in which fat transfer can be used for subsequent contour correction.¹ Although an oncologically permissible environment for fat grafting may exist when residual disease is resected and appropriate adjuvant therapy administered, increased recurrence risk may exist when fat is transferred in the presence of residual breast tissue.^{46–48} The European Institute of Oncology reported a locoregional recurrence rate of 0.4 percent per year following partial mastectomy among 2784 subjects.⁴⁹ However, a retrospective review of 143 partial mastectomy patients found that fat transfer was associated with a 2.07 percent per year increased rate of locoregional recurrence.⁷ By contrast, and in support of our data, recurrence rates in mastectomy patients after fat transfer increased 1.38 percent per year,⁷ versus 1.1 percent per year in a historical control group of 677 patients who did not receive fat transfer.⁵⁰ In the absence of a prospective trial, the authors of this multicenter retrospective review of 646 fat transfer patients recommended a cautious oncologic follow-up protocol.⁷

Fat transfer may be associated with recurrence after ductal carcinoma in situ, which we did not include in our study population. In a matched cohort study of 321 patients, a significantly higher incidence of local and locoregional recurrence after fat transfer was found when analysis was limited to patients with in situ disease.⁶ A subsequent study limited to 59 patients with in situ disease who received fat transfer, and a matched cohort of 118 who did not, revealed a higher rate of locoregional recurrence in the fat transfer group (hazard ratio, 4.5; 95 percent CI, 1.1 to 18.2).⁸ Patients younger than 50 years, with high-grade neoplasia and a Ki-67 value greater than or equal to 14, were also associated with an increased rate of recurrence after fat transfer in this study.

In the retrospective series by Petit et al., fat transfer was performed in 108 patients with in situ disease, with a locoregional recurrence rate of 2.33 percent per year versus 1.44 percent per year in 405 patients with invasive carcinomas.⁷ The molecular signature of the epithelial component of the tumor microenvironment that regulates extracellular matrix remodeling differs between ductal carcinoma in situ and invasive ductal carcinoma.⁵¹ Moreover, subtypes of ductal carcinoma in situ can be differentiated by unique molecular signatures expressed by their fibroblasts and vascular and inflammatory stromal cells.^{52,53} Recognizing these differences, Petit and colleagues⁸ postulated that with fewer genetic perturbations than invasive carcinomas, intraepithelial neoplasias may be more efficient at responding to the stromal signaling that leads to malignant degeneration. Recently, this group reanalyzed their ductal carcinoma in situ study population over a longer period, to increase the number of recurrence events evaluated.²⁶ Relative to controls, the recurrence rate was not higher in patients receiving fat transfer following mastectomy ($p = 0.56$). It was somewhat higher in patients grafted following partial mastectomy, but the difference was not statistically significant ($p = 0.20$). In addition, Gale and colleagues did not report an increased rate of recurrence following fat transfer in ductal carcinoma in situ patients.⁵⁴ Fat transfer in these patients, however, was restricted to patients with clear margins and was delayed for 54 months after resection. By contrast, Petit et al. waited a mean of 25 months and had positive or close margins in 42 percent of fat-transferred patients but only 20 percent of controls, and thus the increased risk of recurrence may have been related to margin status.⁸ Gale et al. suggested that early fat transfer (≤ 2 years) after tumor resection may increase the impact of fat transfer on recurrence rates.⁵⁴ Kronowitz et al.—who also support the oncologic safety of fat transfer in their study—suggest that lumpectomy accompanied by intraoperative radiation therapy only may have also impacted recurrence rates in the study by Petit et al.²²

Evaluation of the oncologic safety of fat transfer to facilitate breast reconstruction is challenging because recurrence of local or regional breast cancer after mastectomy is a relatively rare outcome,⁵⁵ many years of follow-up are necessary to evaluate recurrence,^{56,57} and fat transfer has become common only recently. Thus, we used a retrospective case cohort approach. Our study was sufficiently powered to detect a risk ratio of 2 or greater, in terms of association between fat

transfer and breast cancer recurrence. Moving forward, we intend to continue to study cancer recurrence rates after fat transfer with a larger sample size and longer follow-up. This will improve on the risk ratio of 2 used in this study, and will allow us to comment on the impact of other factors on recurrence such as volume transferred and fat transfer techniques used in a statistically meaningful way. Although a prospective, randomized, controlled trial would be favorable over a retrospective analysis, it would be impractical to ask patients to agree to a control arm of no treatment for contour deformities. Finding an alternative control arm, such as temporary fillers or tissue rearrangement, would also be challenging. A prospective cohort study would also be useful but would require a substantially larger sample size and would not produce evidence for 5 to 10 years. Finally, a study using existing large administrative or clinical data sets is not feasible because fat transfer does not yet have a unique billing code and is not routinely recorded in cancer registries. To address an urgent need for evidence on the safety of fat transfer to the breast,⁵⁸ we used a case-cohort study design, which is appropriate for assessing the probability of rare outcomes in a more timely fashion than a prospective cohort study.

The retrospective nature of this study is a limitation. In addition, we did not adjust for differences among fat transfer techniques because of a lack of consensus, nuanced technique differences not captured by retrospective review, and insufficient power to evaluate the impact of different fat transfer techniques on recurrence. Still, various methods of fat harvest and processing may affect adipocyte viability and stem cell fraction.^{59–62} Although optimization of the adipose-derived stem cell–rich stromal vascular fraction of lipoaspirate may favor improved graft retention,⁴⁵ it may also increase the risk of exposure of adipose-derived stem cells to occult, residual tumor stroma.⁴⁵ All patients were treated at high-volume cancer centers and likely had access to timely cancer treatment and appropriate administration of adjuvant therapy.^{48,63} Thus, these results assume guideline-concordant care and may not be generalizable to all breast cancer patients. In addition, some of these patients may have sought a cancer center for their initial therapy but eluded detection by the institutional tumor registry for recurrences treated elsewhere. We assessed only patients undergoing mastectomy and immediate reconstruction, a population that tends to be healthier, has greater economic

resources, and has more favorable tumors than breast cancer patients overall.^{64–66} Although we adjusted for some clinical variables, we did not adjust for health status or social factors. Future studies should include patients who have fat transfer after delayed reconstruction and after breast conservation therapy. As fat transfer to the breast becomes more common and acquires more indications, such studies ought to become more feasible.

CONCLUSIONS

Fat transfer was not associated with a higher probability of recurrence in this multisite population of local and regional breast cancer patients treated with mastectomy and reconstruction. Although the precision of the study was somewhat limited, it provides evidence that fat transfer does not increase the probability of invasive breast cancer recurrence by a factor of at least 2.0. Future studies of a larger sample of immediate reconstruction patients, and studies of fat transfer after delayed reconstruction or breast conservation therapy, are warranted.

Terence M. Myckatyn, M.D.

Plastic and Reconstructive Surgery

Washington University School of Medicine in St. Louis

Saint Louis, Mo. 63110

myckatyn@wudosis.wustl.edu

ACKNOWLEDGMENTS

This work was supported by The Plastic Surgery Foundation. Clara Lee, M.D., M.P.P., was supported by National Cancer Institute grant 1K07CA154850-01A1. Evan Busch, Ph.D., was supported in part by National Cancer Institute grant 5T32CA009001. The authors wish to thank Lillian Blizard (University of North Carolina) for technical support, Katie Sommers (The Plastic Surgery Foundation) for project coordination support, Colleen Kilbourne-Glynn (Washington University School of Medicine in St. Louis), Tracie Guthrie (Washington University School of Medicine in St. Louis), Sara Baalman (Washington University School of Medicine in St. Louis), Meghan Lee (Memorial Sloan Kettering Cancer Center), Debbie Crawford (M. D. Anderson Cancer Center) and Kelly Wall (M. D. Anderson Cancer Center) for assistance with data collection.

REFERENCES

1. Spear SL. Fat for breast: Where are we? *Plast Reconstr Surg.* 2008;122:983–984.
2. American Society of Plastic Surgeons. 2014 plastic surgery statistics. Available at: <https://www.plasticsurgery.org/news/>

- plastic-surgery-statistics?sub=2014+Plastic+Surgery+Statistics. Accessed April 15, 2015.
- Kling RE, Mehrara BJ, Pusic AL, et al. Trends in autologous fat grafting to the breast: A national survey of the American Society of Plastic Surgeons. *Plast Reconstr Surg*. 2013;132:35–46.
 - Bertolini F, Lohsiriwat V, Petit JY, Kolonin MG. Adipose tissue cells, lipotransfer and cancer: A challenge for scientists, oncologists and surgeons. *Biochim Biophys Acta* 2012;1826:209–214.
 - Bertolini F, Petit JY, Kolonin MG. Stem cells from adipose tissue and breast cancer: Hype, risks and hope. *Br J Cancer* 2015;112:419–423.
 - Petit JY, Botteri E, Lohsiriwat V, et al. Locoregional recurrence risk after lipofilling in breast cancer patients. *Ann Oncol*. 2012;23:582–588.
 - Petit JY, Lohsiriwat V, Clough KB, et al. The oncologic outcome and immediate surgical complications of lipofilling in breast cancer patients: A multicenter study—Milan-Paris-Lyon experience of 646 lipofilling procedures. *Plast Reconstr Surg*. 2011;128:341–346.
 - Petit JY, Rietjens M, Botteri E, et al. Evaluation of fat grafting safety in patients with intraepithelial neoplasia: A matched-cohort study. *Ann Oncol*. 2013;24:1479–1484.
 - Pearl RA, Leedham SJ, Pacifico MD. The safety of autologous fat transfer in breast cancer: Lessons from stem cell biology. *J Plast Reconstr Aesthet Surg*. 2012;65:283–288.
 - Zhang Y, Daquinag A, Traktuev DO, et al. White adipose tissue cells are recruited by experimental tumors and promote cancer progression in mouse models. *Cancer Res*. 2009;69:5259–5266.
 - Manabe Y, Toda S, Miyazaki K, Sugihara H. Mature adipocytes, but not preadipocytes, promote the growth of breast carcinoma cells in collagen gel matrix culture through cancer-stromal cell interactions. *J Pathol*. 2003;201:221–228.
 - Schäffler A, Schölmerich J, Buechler C. Mechanisms of disease: Adipokines and breast cancer. Endocrine and paracrine mechanisms that connect adiposity and breast cancer. *Nat Clin Pract Endocrinol Metab*. 2007;3:345–354.
 - Iyengar P, Espina V, Williams TW, et al. Adipocyte-derived collagen VI affects early mammary tumor progression in vivo, demonstrating a critical interaction in the tumor/stroma microenvironment. *J Clin Invest*. 2005;115:1163–1176.
 - Lohsiriwat V, Curigliano G, Rietjens M, Goldhirsch A, Petit JY. Autologous fat transplantation in patients with breast cancer: “Silencing” or “fueling” cancer recurrence? *Breast* 2011;20:351–357.
 - Zimmerlin L, Donnenberg AD, Rubin JP, Basse P, Landreneau RJ, Donnenberg VS. Regenerative therapy and cancer: In vitro and in vivo studies of the interaction between adipose-derived stem cells and breast cancer cells from clinical isolates. *Tissue Eng Part A* 2011;17:93–106.
 - Martin-Padura I, Gregato G, Marighetti P, et al. The white adipose tissue used in lipotransfer procedures is a rich reservoir of CD34+ progenitors able to promote cancer progression. *Cancer Res*. 2012;72:325–334.
 - Orecchioni S, Gregato G, Martin-Padura I, et al. Complementary populations of human adipose CD34+ progenitor cells promote growth, angiogenesis, and metastasis of breast cancer. *Cancer Res*. 2013;73:5880–5891.
 - Direkze NC, Jeffery R, Hodivala-Dilke K, et al. Bone marrow-derived stromal cells express lineage-related messenger RNA species. *Cancer Res*. 2006;66:1265–1269.
 - Kolonin MG, Evans KW, Mani SA, Gomer RH. Alternative origins of stroma in normal organs and disease. *Stem Cell Res*. 2012;8:312–323.
 - Karnoub AE, Dash AB, Vo AP, et al. Mesenchymal stem cells within tumour stroma promote breast cancer metastasis. *Nature* 2007;449:557–563.
 - Krastev TK, Jonasse Y, Kon M. Oncological safety of autologous lipoaspirate grafting in breast cancer patients: A systematic review. *Ann Surg Oncol*. 2013;20:111–119.
 - Kronowitz SJ, Mandujano CC, Liu J, et al. Lipofilling of the breast does not increase the risk of recurrence of breast cancer: A matched controlled study. *Plast Reconstr Surg*. 2016;137:385–393.
 - American Society of Plastic Surgeons. Post-mastectomy fat graft/fat transfer ASPS guiding principles. Available at: <http://www.plasticsurgery.org/Documents/Health-Policy/Principles/principle-2015-post-mastectomy-fat-grafting.pdf>. Accessed August 15, 2015.
 - Cai J, Zeng D. Sample size/power calculation for case-cohort studies. *Biometrics* 2004;60:1015–1024.
 - R Foundation. A language and environment for statistical computing: R Foundation for Statistical Computing. Available at: <http://www.r-project.org/>. Accessed August 6, 2015.
 - Petit JY, Maisonneuve P, Rotmensz N, et al. Safety of lipofilling in patients with breast cancer. *Clin Plast Surg*. 2015;42:339–344, viii.
 - Khoury RK, Smit JM, Cardoso E, et al. Percutaneous aponeurotomy and lipofilling: A regenerative alternative to flap reconstruction? *Plast Reconstr Surg*. 2013;132:1280–1290.
 - Cleary MP, Grossmann ME, Ray A. Effect of obesity on breast cancer development. *Vet Pathol*. 2010;47:202–213.
 - Dirat B, Bochet L, Escourrou G, Valet P, Muller C. Unraveling the obesity and breast cancer links: A role for cancer-associated adipocytes? *Endocr Dev*. 2010;19:45–52.
 - Hede K. Fat may fuel breast cancer growth. *J Natl Cancer Inst*. 2008;100:298–299.
 - Rose DP, Haffner SM, Baillargeon J. Adiposity, the metabolic syndrome, and breast cancer in African-American and white American women. *Endocr Rev*. 2007;28:763–777.
 - Zhang Y, Bellows CF, Kolonin MG. Adipose tissue-derived progenitor cells and cancer. *World J Stem Cells* 2010;2:103–113.
 - Zhang Y, Daquinag AC, Amaya-Manzanares F, Sirin O, Tseng C, Kolonin MG. Stromal progenitor cells from endogenous adipose tissue contribute to pericytes and adipocytes that populate the tumor microenvironment. *Cancer Res*. 2012;72:5198–5208.
 - Kidd S, Spaeth E, Watson K, et al. Origins of the tumor microenvironment: Quantitative assessment of adipose-derived and bone marrow-derived stroma. *PLoS One* 2012;7:e30563.
 - Li H, Zimmerlin L, Marra KG, Donnenberg VS, Donnenberg AD, Rubin JP. Adipogenic potential of adipose stem cell subpopulations. *Plast Reconstr Surg*. 2011;128:663–672.
 - Ray A, Cleary MP. Leptin as a potential therapeutic target for breast cancer prevention and treatment. *Expert Opin Ther Targets* 2010;14:443–451.
 - Grossmann ME, Ray A, Nkhata KJ, et al. Obesity and breast cancer: Status of leptin and adiponectin in pathological processes. *Cancer Metastasis Rev*. 2010;29:641–653.
 - Núñez NP, Carpenter CL, Perkins SN, et al. Extreme obesity reduces bone mineral density: Complementary evidence from mice and women. *Obesity (Silver Spring)* 2007;15:1980–1987.
 - Taliaferro-Smith L, Nagalingam A, Knight BB, Oberlick E, Saxena NK, Sharma D. Integral role of PTP1B in adiponectin-mediated inhibition of oncogenic actions of leptin in breast carcinogenesis. *Neoplasia* 2013;15:23–38.
 - Dirat B, Bochet L, Dabek M, et al. Cancer-associated adipocytes exhibit an activated phenotype and contribute to breast cancer invasion. *Cancer Res*. 2011;71:2455–2465.

41. Chandler EM, Seo BR, Califano JP, et al. Implanted adipose progenitor cells as physicochemical regulators of breast cancer. *Proc Natl Acad Sci USA* 2012;109:9786–9791.
42. Eterno V, Zambelli A, Pavesi L, et al. Adipose-derived mesenchymal stem cells (ASCs) may favour breast cancer recurrence via HGF/c-Met signaling. *Oncotarget* 2014;5:613–633.
43. Rowan BG, Gimble JM, Sheng M, et al. Human adipose tissue-derived stromal/stem cells promote migration and early metastasis of triple negative breast cancer xenografts. *PLoS One* 2014;9:e89595.
44. Phillips BJ, Grahovac TL, Valentin JE, et al. Prevalence of endogenous CD34+ adipose stem cells predicts human fat graft retention in a xenograft model. *Plast Reconstr Surg*. 2013;132:845–858.
45. Meyer J, Salamon A, Herzmann N, et al. Isolation and differentiation potential of human mesenchymal stem cells from adipose tissue harvested by water jet-assisted liposuction. *Aesthet Surg J*. 2015;35:1030–1039.
46. Vicini FA, Eberlein TJ, Connolly JL, et al. The optimal extent of resection for patients with stages I or II breast cancer treated with conservative surgery and radiotherapy. *Ann Surg*. 1991;214:200–204; discussion 204.
47. McCahill LE, Single RM, Aiello Bowles EJ, et al. Variability in reexcision following breast conservation surgery. *JAMA* 2012;307:467–475.
48. Morrow M, Harris JR, Schnitt SJ. Surgical margins in lumpectomy for breast cancer: Bigger is not better. *N Engl J Med*. 2012;367:79–82.
49. Botteri E, Bagnardi V, Rotmensz N, et al. Analysis of local and regional recurrences in breast cancer after conservative surgery. *Ann Oncol*. 2010;21:723–728.
50. Petit JY, Gentilini O, Rotmensz N, et al. Oncological results of immediate breast reconstruction: Long term follow-up of a large series at a single institution. *Breast Cancer Res Treat*. 2008;112:545–549.
51. Vargias AC, McCart Reed AE, Waddell N, et al. Gene expression profiling of tumour epithelial and stromal compartments during breast cancer progression. *Breast Cancer Res Treat*. 2012;135:153–165.
52. Sharma M, Beck AH, Webster JA, et al. Analysis of stromal signatures in the tumor microenvironment of ductal carcinoma in situ. *Breast Cancer Res Treat*. 2010;123:397–404.
53. Orimo A, Gupta PB, Sgroi DC, et al. Stromal fibroblasts present in invasive human breast carcinomas promote tumor growth and angiogenesis through elevated SDF-1/CXCL12 secretion. *Cell* 2005;121:335–348.
54. Gale KL, Rakha EA, Ball G, Tan VK, McCulley SJ, Macmillan RD. A case-controlled study of the oncologic safety of fat grafting. *Plast Reconstr Surg*. 2015;135:1263–1275.
55. Bouganim N, Tsvetkova E, Clemons M, Amir E. Evolution of sites of recurrence after early breast cancer over the last 20 years: Implications for patient care and future research. *Breast Cancer Res Treat*. 2013;139:603–606.
56. Mansi JL, Gogas H, Bliss JM, Gazet JC, Berger U, Coombes RC. Outcome of primary-breast-cancer patients with micrometastases: A long-term follow-up study. *Lancet* 1999;354:197–202.
57. Braun S, Vogl FD, Naume B, et al. A pooled analysis of bone marrow micrometastasis in breast cancer. *N Engl J Med*. 2005;353:793–802.
58. Gutowski KA; ASPS Fat Graft Task Force. Current applications and safety of autologous fat grafts: A report of the ASPS fat graft task force. *Plast Reconstr Surg*. 2009;124:272–280.
59. Coleman SR. Structural fat grafting: More than a permanent filler. *Plast Reconstr Surg*. 2006;118(Suppl):108S–120S.
60. Khouri R, Del Vecchio D. Breast reconstruction and augmentation using pre-expansion and autologous fat transplantation. *Clin Plast Surg*. 2009;36:269–280, viii.
61. Khouri RK, Rigotti G, Cardoso E, Khouri RK Jr, Biggs TM. Megavolume autologous fat transfer: Part II. Practice and techniques. *Plast Reconstr Surg*. 2014;133:1369–1377.
62. Gir P, Brown SA, Oni G, Kashefi N, Mojallal A, Rohrich RJ. Fat grafting: Evidence-based review on autologous fat harvesting, processing, reinjection, and storage. *Plast Reconstr Surg*. 2012;130:249–258.
63. Mamounas EP, Tang G, Fisher B, et al. Association between the 21-gene recurrence score assay and risk of locoregional recurrence in node-negative, estrogen receptor-positive breast cancer: Results from NSABP B-14 and NSABP B-20. *J Clin Oncol*. 2010;28:1677–1683.
64. Wilkins EG, Alderman AK. Breast reconstruction practices in North America: Current trends and future priorities. *Semin Plast Surg*. 2004;18:149–155.
65. Joslyn SA. Patterns of care for immediate and early delayed breast reconstruction following mastectomy. *Plast Reconstr Surg*. 2005;115:1289–1296.
66. Christian CK, Niland J, Edge SB, et al. A multi-institutional analysis of the socioeconomic determinants of breast reconstruction: A study of the National Comprehensive Cancer Network. *Ann Surg*. 2006;243:241–249.